

PHASE-II[®]

A New Generation MSO Adjuvant Oil

An emulsifiable concentrate containing 842 g/l (95,2 % w/w) esterified rapeseed oil



DE SANGOSSE

De Sangosse Ltd
Swaffham Bulbeck, Cambridge CB25 0LU
Tel: 01223 811215 Fax: 01223 810020
e-mail: Info@desangosse.co.uk web: www.desangosse.co.uk
Distributed in Ireland by
T.P. Whelehan Son & Co Ltd, Finglas Dublin 11, tel: 01 8058600

FOR USE ONLY AS AN
AGRICULTURAL/HORTICULTURAL ADJUVANT OIL

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

PROTECT FROM FROST

ADJ No 0622 PCS 92147

5 Litres e

PCS: 92147

ADJ NO: 0622

RESERVE DE TEXTE



**STATUTORY CONDITIONS RELATING TO USE
FOR USE ONLY AS AN AGRICULTURAL/
HORTICULTURAL ADJUVANT OIL**
Maximum Concentration: 1.78% Spray Solution
ADJ No 0622 PCS 92147
READ ALL PRECAUTIONS BEFORE USE



PRECAUTIONS

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

OPERATOR PROTECTION

This product contains no hazardous substances under the directive 1999/45/EC

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS), SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the concentrate. However engineering controls may replace personal protective equipment if a COSHH assessment shows that they provide an equal or higher standard of protection.

DO NOT BREATHE SPRAY/MISTS. WASH HANDS AND EXPOSED SKIN before meals and after work.

TO AVOID RISKS TO MAN AND THE ENVIRONMENT, COMPLY WITH THE INSTRUCTIONS FOR USE.

Operators must follow all operator protection instructions for the pesticide. Where the personal protective equipment specified on the adjuvant label is different to that specified on the pesticide label, the highest level of protective equipment must be worn.

ENVIRONMENTAL PROTECTION

DO NOT CONTAMINATE SURFACE WATER OR DITCHES with chemical or used container. **DO NOT EMPTY INTO DRAINS;** dispose of this material and its container in a safe way.

STORAGE AND DISPOSAL
KEEP OUT OF REACH OF CHILDREN.

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS.

WASH OUT CONTAINER THOROUGHLY, and dispose of safely.

DO NOT RE-USE CONTAINER for any purpose.

CONDITIONS

PRECAUTIONS



DIRECTIONS FOR USE

PHASE-II may be used at a maximum concentration of 1.78% spray solution (maximum 2 litres per hectare) on the following crops, up to the growth stages specified, with all approved pesticides applied up to their full approved rate.



Crop	Growth Stages before which Phase-II may be used with approved pesticides
Cereals	Up to and including GS 52 (1/4 inflorescence)
Oilseed rape	Up to and including 10% potential pods
Broad bean, dwarf French bean, field bean, runner bean	Up to and including first pod set
Combining pea & vining pea	Up to and including flat pod (non-edible pods only)
Edible podded pea	Up to and including pod set
Linseed	Up to and including 10% capsules formed
Forage Maize	Up to and including tip of tassel visible
Sugar beet	Up to and including 6 fully expanded leaves
Potato	Up to and including tuber initiation
Brussels sprout	Up to and including lateral buds begin to develop
Cauliflower	Up to and including cauliflower heads begin to form; width of growing tip up to 1 cm ²

In addition to the above conditions PHASE-II is authorised for use with all approved pesticides on all edible crops provided that the statutory conditions of use for the pesticide are followed and that the pesticide is not used at more than 50% of the maximum approved rate for that application.

PHASE-II is also authorised for use with all approved pesticides on all non-edible crops provided that the statutory conditions of use for the pesticide are followed.

PHASE-II may also be used to reduce drift of sprays where this is a concern.

For Specific Recommendations contact De Sangosse Ltd or its appointed Distributors.

DIRECTIONS

RESERVE DE TEXTE



RATES OF USE

PHASE-II is authorised for use up to a maximum concentration of 1.78% dilution of the final water volume or a maximum dose of 2 litres per hectare. In most situations the normal recommended rate for PHASE-II is 0.5% e.g.

Water Volume	Required Quantity of Phase-II at 0.5%
100 litres	500 ml
200 litres	1,000 ml

Rates of PHASE-II can vary depending on the partner pesticide. For specific rate of use recommendations please consult your De Sangosse Ltd. Distributor.



MIXING INSTRUCTIONS

Three quarter fill the spray tank with water and commence agitation. Add the required quantity of PHASE-II then add the recommended quantity of pesticide(s) as directed. Continue agitation while topping up the tank and during spraying.

CONDITIONS OF SUPPLY

All goods supplied by us are of high grade and we believe them to be suitable, but as we cannot exercise control over their storage, mixing or use, all conditions and warranties, statutory or otherwise as to the quality or fitness for any purpose of our goods are excluded and no responsibility will be accepted by us for any damage or injury whatsoever arising from their storage, handling, application or use.

PHASE-II is a Trademark of De Sangosse Ltd.

RATES

MIXING

SUPPLY