



FEED, WEED & MOSS KILLER



MAPP No. 19669

4 Plication

BALANCED NPK
ANALYSIS FOR
USE THROUGHOUT
THE GROWING
SEASON.





Scan me to view data sheet online.



STORE IN A COOL & DRY PLACE

5kg



MAPP No. 19669

INSTRUCTIONS FOR USE

WHEN TO USE:



Feed, Weed & Moss Killer can be used between May and August when grass, moss and weeds are actively growing, and soil is moist. Do not apply during drought or freezing conditions. Granules and dust should not be allowed to come in to contact with flowers, shrubs, fruit and vegetable plants.

Apply only to established turf. Do not use on new grass for six months. Grass should not be cut 3 days before treatment or until at least 4 days after treatment. Do not apply when heavy rain is forecast, but if no rain falls within 48 hours, water the product into the lawn.

HOW TO USE:



APPLICATION BY SPREADER:

Mark out areas to be treated. Apply evenly at 32 grams per square metre over the marked area with a suitable calibrated granulated applicator.



APPLICATION BY SHAKER PACK OR BY HAND:

Mark out areas to be treated. Apply evenly at 32 grams per square metre over the marked area with shaker pack or by hand. It is good practice to wear gloves when using this product although this is not a requirement. Do not apply in windy conditions. Keep off skin and away from eyes. Avoid walking on treated areas until it has rained, or product has been watered in. If spilt on paving, concrete, clothing etc. brush off immediately to avoid discolouration. If no rain falls within 48 hours of treatment, irrigate the treated area.



FURTHER INFORMATION:

The moss will blacken as it dies. Any discolouration of the turf is temporary. Do not re-seed or turf for at least 6 weeks after application. The first four mowings should not be used to mulch cultivated plants unless composted for at least six months. Feed, Weed & Moss Killer must not be mixed or used with other plant protection products.

Dead Moss can be removed by raking or scarifying 7-14 days after treatment. Pre-treatment may be necessary for heavy moss infestation, or if moss returns. Do not apply more than 2 times in any one season. A 40 day interval is required between applications.



KEEP CHILDREN AND PETS AWAY FROM TREATED AREAS UNTIL **GRANULES HAVE DISSOLVED.**

Feed, Weed & Moss Killer is effective against broad leaves weeds of the following species:

- **Common Dandelion** (Taraxacum officinale)
- Creeping Buttercup (Ranunculus repens)
- Creeping Cinquefoil (Potentilla reptans)
- Daisy (Bellis perennis)
- Greater Plantain (Plantago major)
- Lesser Trefoil (Trifolium Dubium)

- Mouse-ear Chickweed (Cerastium fontanum)
- Ribwort, Plantain (Plantago lanceolata)
- Selfheal (Prunella Vulgaris)
- Thyme-leaved Speedwell (Veronica serpyllifolia)
- White Clover (Trifolium repens)
- Birds Foot Trefoil (Lotus Corniculatus)

Large or perennial weeds may regrow. Persistent weeds may require repeated treatments once the new growth appears.

CONTROL TECHNIOUES:

The key to reducing moss and weeds is to have an effective all year-round management strategy incorporating Feed, Weed & Moss Killer.

The following non-chemical techniques can be used to help reduce moss and weeds:

- Ensure the turf is healthy, weed, disease and pest free.
- This can help by fertilisation and regular maintenance.
- Soil aerating of the lawn to help to encourage the free passage of water & nutrients through the soil.
- Raking / scarifying to remove dead mosses and removal of fallen leaf debris to prevent the build-up of fibrous material which encourages weed and moss growth.

If all these cultural management practices are followed in harmony with applications of Feed, Weed & Moss Killer than the reduction of mosses and weeds can be successfully achieved.

Scan me data sheet online.





STORE IN A COOL & DRY PLACE

Agrigem Ltd, Riverside Enterprise Park, Gem House, Skellingthorpe Road, Saxilby LN1 2FU Tel: 01522 246491





MAPP No. 19669

FEED, WEED & MOSS KILLER

Causes serious eye irritation.

Harmful to aquatic life with long lasting effects.

If medical advice is needed, have product container or label at hand.



Wash hands thoroughly after handling.

Wear protective gloves / protective clothing /

eye protection / face protection.

If eye irritation persists, seek medical attention.

Collect spillage.

Dispose of contents at a household waste recycling centre as hazardous waste, except for empty container which can be deposed of by recycling. Contact your local council for details.

Contains 2, 4-D. May produce an allergic reaction.

To avoid risk to human health and the environment. comply with the instructions for use.

SAFETY INSTRUCTIONS:

PROTECTION DURING USE:

WASH HANDS AND EXPOSED SKIN AFTER USE / HANDLING.

If swallowed seek medical attention immediately and show this packaging or label.

ENVIRONMENT PROTECTION:

Do not contaminate water with the product or packaging. Do not empty into drains.

STORAGE AND DISPOSAL:

Store in a cool dry place.

KEEP PRODUCT IN ORIGINAL PACKAGING in a safe place.

KEEP PRODUCT AWAY FROM CHILDREN AND PETS.

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEED.

COMPOUND FERTILISER WITH IRON

10 - 2 - 1.7 UK

Nitrogen (N)

10%

Total Phosphorous Pentoxide (P.O.) Of which is soluble in water.

2.0% (0.9%P)

Total Potassium Oxide (K_o O)

1.1% (O.5%P)

1.7% (1.4%K)

Iron (Fe) soluble in water

8%

Scan me to view data sheet online.



Treats 160m²

STORE IN A COOL & DRY PLACE

Agrigem Ltd, Riverside Enterprise Park, Gem House, Skellingthorpe Road, Saxilby LN1 2FU Tel: 01522 246491

