

INTER TEBLOXY F

Inter Tebloxy is a contact and systemic fungicide containing 200g/l (18.2%w/w) tebuconazole and 100 g/l (9.10%w/w) trifloxystrobin.

Inter Tebloxy has preventative and early curative action against Red Thread (*Laetisaria fuciformis*), Fusarium Patch (*Microdochium nivale*), Anthracnose (*Colletotrichum cereale*) Leaf Spot (*Drechslera* spp.), Dollar Spot (*Sclerotinia homoeocarpa*) and Rust (*Puccinia* spp) at any time of year in managed amenity turf.

Registration Number: MAPP 15412



DANGEROUS FOR THE ENVIRONMENT



HARMFUL

VERY TOXIC TO AQUATIC ORGANISMS, MAY CAUSE LONG-TERM ADVERSE EFFECTS IN THE AQUATIC ENVIRONMENT.

POSSIBLE RISK OF HARM TO THE UNBORN CHILD. IRRITATING TO THE RESPIRATORY SYSTEM.

Do not breathe spray.

This material and its container must be disposed of in a safe way.

Wear suitable protective clothing and gloves.

Use appropriate containment to avoid environmental contamination.

May produce an allergic reaction. Contains trifloxystrobin.

To avoid risks to man and the environment comply with the instructions for use.

National Poisons Information Service: 0844 892 0111



This label is compliant with the CPA Voluntary Initiative Guidance

1 Litre e

SAFETY PRECAUTIONS

Operator Protection

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment : WEAR SUITABLE PROTECTIVE CLOTHING AND SUITABLE PROTECTIVE GLOVES when handling the concentrate, applying by hand-held equipment or handling contaminated surfaces. However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection. DO NOT BREATHE SPRAY. WHEN USING DO NOT EAT, DRINK OR SMOKE. WASH ANY CONTAMINATION from eyes immediately. IF YOU FEEL UNWELL, seek medical advice (show label where possible). WASH HANDS AND EXPOSED SKIN before meals and after work.

Environmental Protection

DO NOT CONTAMINATE SURFACE WATERS OR DITCHES with chemical or used container.

Storage and Disposal

KEEP OUT OF REACH OF CHILDREN. KEEP IN ORIGINAL CONTAINER tightly closed in a safe place. WASH OUT CONTAINER THOROUGHLY, empty washings into spray tank and dispose of safely. DO NOT RE-USE CONTAINER for any purpose. KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS.

Read the label before use. Using this product in a manner that is inconsistent with the label may be an offence. Follow the code of practice for using plant protection products.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

Inter Tebloxy is a fungicide which can be used on areas of established turf, for example golf courses, sports fields, municipal parkland, residential lawns and cemeteries.

General

Inter Tebloxy is a combination of two active ingredients (tebuconazole and trifloxystrobin). These two actives are from two different fungicide groups; a triazole (tebuconazole) and a strobilurin (trifloxystrobin). It is designed to control Fusarium and reduce the incidence of a range of other turf diseases in managed amenity turf.

Important

Do not make applications of Inter Tebloxy in drought conditions or on to frozen turf.

Rate of application

In order to control Fusarium, Red Thread and Dollar Spot apply at 1 litre of Inter Tebloxy per hectare in a minimum of 400-500 litres of water. At this rate some control of Anthracnose and reduction of rust may be achieved. Apply by either a vehicle mounted hydraulic sprayer or knapsack sprayer as described below:

Application Equipment			
Vehicle-mounted Hydraulic Sprayer		Knapsack Sprayer	
Rate per hectare	Water per hectare	Rate/100 m ²	Water/100m ²
1 Litre	400 – 500 litres	10 mls	4 – 5 litres
		Rate/500 m ²	Water/500m ²
		50 mls	20 – 25 litres

Optimum results will be achieved if Inter Tebloxy is applied at the very first signs of Fusarium Infection, or at a very early stage in the development of the disease. Inter Tebloxy can be applied when the grass is actively growing or when it is dormant, as the product has both curative and protective properties. During the mowing season apply after cutting and do not mow again until at least 48 hours after treatment. This 48 hour period allows the product to move systematically throughout the plant. Apply a maximum of twice per year. Additionally allow a minimum of 28 days between application of Inter Tebloxy. Adopt fungicide management programmes which combine fungicides with different modes of action. Try to avoid spraying Inter Tebloxy in sequence.

Resistance Management

Always try to avoid the build up of disease resistance. Repeated use of the same active or product from the same fungicide group may result in resistance occurring. Always alternate different fungicides if possible in a programmed approach.

Mixing

Half fill the spray tank with clean water and commence agitation. Add the required quantity of Inter Tebloxy and continue filling the tank. Maintain agitation during spraying. Do not allow the mixture to stand for long periods before use. Always ensure that the sprayer is clean, in good working order and calibrated to the sprayer manufacturer's instructions.

Iticon N.V. Werfkaai 14, B-8380, Zeebrugge, Belgium
Tel: 0032 50 550 777 Fax:0032 50 550 716